

CLASSIFYING p -GROUPS VIA THEIR MULTIPLIER

PEYMAN NIROOMAND

ABSTRACT. The author in (On the order of Schur multiplier of non-abelian p -groups. J. Algebra (2009).322: 4479–4482) showed that for any p -group G of order p^n there exists a nonnegative integer $s(G)$ such that the order of Schur multiplier of G is equal to $p^{\frac{1}{2}(n-1)(n-2)+1-s(G)}$. Furthermore, he characterized the structure of all non-abelian p -groups G when $s(G) = 0$. The present paper is devoted to characterization of all p -groups when $s(G) = 2$.

The concept of Schur multiplier, $\mathcal{M}(G)$, have been studied by several authors, initiated by Schur in 1904. It is known that the order of Schur multiplier of a given finite p -group of order p^n is equal to $p^{\frac{1}{2}n(n-1)-t(G)}$ for some $t(G) \geq 0$ by a result of Green [5]. It is of interest to know which p -groups have the Schur multiplier of order $p^{\frac{1}{2}n(n-1)-t(G)}$, when $t(G)$ is in hand.

Historically, there are several papers trying to characterize the structure of G by just the order of its Schur multiplier. In [1] and [13], Berkovich and Zhou classified the structure of G when $t(G) = 0, 1$ and 2 , respectively.

Later, Ellis in [2] showed that having a new upper bound on the order of Schur multiplier of groups reduces characterization process of structure of G . He reformulated the upper bound due to Gaschütz at. al. [4] and classified with a quite way to that of [1, 13] the structure of G when $t(G) = 3$.

The result of [9] shows that there exists a nonnegative integer $s(G)$ such that $|\mathcal{M}(G)| = p^{\frac{1}{2}(n-1)(n-2)+1-s(G)}$ which is a reduction of Green's bound for any given non-abelian p -group G of order p^n . One can check that the structure of G can be characterized by using [9, Main Theorem], when $t(G) = 1, 2, 3$. Moreover, characterizing non-abelian p -groups by $s(G)$ can be significant since for instance the result of [9] and [12] emphasize that the number of groups with a fixed $s(G)$ is more than that with fixed $t(G)$. Also the results of [10] and [11] show handling the p -groups characterized by $s(G) = 0, 1$ may be caused to characterize the structure of G by $t(G)$.

In the present paper, we intend to classify the structure of all non-abelian p -groups when $s(G) = 2$.

Throughout this paper we use the following notations.

Q_8 : quaternion group of order 8,

D_8 : dihedral group of order 8,

E_1 : extra special p -group of order p^3 and exponent p ,

E_2 : extra special p -group of order p^3 and exponent p^2 ($p \neq 2$),

$\mathbb{Z}_{p^n}^{(m)}$: direct product of m copies of the cyclic group of order p^n ,

G^{ab} : the abelianization of group G ,

$H \cdot K$: the central product of H and K ,

Date: March 4, 2010.

Mathematics Subject Classification 2010. Primary 20D15; Secondary 20E34, 20F18.

$E(m)$: $E \cdot Z(E)$, where E is an extra special p -group and $Z(E)$ is cyclic group of order p^m ($m \geq 2$),

$\Phi(G)$: the Frattini subgroup of group G ,

Also, G has the property $s(G) = 2$ or briefly with $s(G) = 2$ means the order of its Schur multiplier is of order $p^{\frac{1}{2}(n-1)(n-2)-1}$.

The following lemma is a consequence of [9, Main Theorem].

Lemma 1. *There is no p -group with $|G'| \geq p^3$ and $s(G) = 2$.*

Lemma 2. *There is no p -group of order p^n ($n \geq 5$) when G^{ab} is not elementary abelian and $s(G) = 2$.*

Proof. First suppose that $n = 5$. By virtue of [11, Theorem 3.6], the result follows. In case $n \geq 6$, by invoking [9, Lemma 2.3], we have $|\mathcal{M}(G/G')| \leq p^{\frac{1}{2}(n-2)(n-3)}$, and since $G/Z(G)$ is capable the rest of proof is obtain by using [3, Propsition 1]. $\square$

Lemma 3. *Let G be a p -group and $|G'| = p$ or p^2 with $s(G) = 2$. Then $Z(G)$ is of exponent at most p^2 and p , respectively.*

Proof. Taking a cyclic central subgroup K of order p^k ($k \geq 3$) and using [6, Theorem 2.2], we should have

$$\begin{aligned} |\mathcal{M}(G)| &\leq p^{-1}|G/K \otimes K|p^{\frac{1}{2}(n-k)(n-k-1)} \leq p^{n-k-1}p^{\frac{1}{2}(n-k)(n-k-1)} \\ &\leq p^{\frac{1}{2}(n-1)(n-2)-2}, \end{aligned}$$

which is a contradiction. In case $|G'| = p^2$, the result obtained similarly. $\square$

Lemma 1 indicates when G has the property $s(G) = 2$, then $|G'| \leq p^2$. First we suppose that $|G'| = p$.

Theorem 4. *Let G be a p -group with centre of order at most p^2 and G^{ab} be elementary abelian of order p^{n-1} and $s(G) = 2$. Then $G \cong E(2)$, $E_2 \times \mathbb{Z}_p$, Q_8 or H , where H is an extra special p -group of order p^{2m+1} ($m \geq 2$).*

Proof. First assume that $|Z(G)| = p$. Hence $G \cong Q_8$ or $G \cong H$, where H is an extra special p -group of order p^{2m+1} ($m \geq 2$) by a result of [8, Theorem 3.3.6]. Now, assume that $|Z(G)| \geq p^2$. Lemma 3 and assumption show that $Z(G) \cong \mathbb{Z}_p \times \mathbb{Z}_p$ or $\mathbb{Z}_{p^2}$.

In case for which $Z(G)$ is of exponent p , [9, Lemma 2.1] follows that $G \cong H \times \mathbb{Z}_p$. It is easily checked that $H \cong E_2$ by using [8, Theorems 2.2.10 and 3.3.6].

In case $Z(G)$ is of exponent p^2 , since $\Phi(G) = G'$, [7, Theorem 3.1] shows that

$$p^{\frac{1}{2}(n-1)(n-2)} = |\mathcal{M}(G/\Phi(G))| \leq p |\mathcal{M}(G)|,$$

and hence $p^{\frac{1}{2}(n-1)(n-2)-1} \leq |\mathcal{M}(G)|$. On the other hand, Main Theorems of [9] and [12] imply that $|\mathcal{M}(G)| = p^{\frac{1}{2}(n-1)(n-2)-1}$ since $Z(G)$ is cyclic of order p^2 . Moreover, $G \cong E(2)$ by appealing [9, Lemma 2.1], as required. $\square$

Theorem 5. *Let G be a p -group, G^{ab} be elementary abelian, $|G'| = p$ and $|Z(G)| \geq p^3$ be of exponent p^2 . Then $G = E(2) \times Z$, where Z is an elementary abelian p -group.*

Proof. It is known that $G = H \cdot Z(G)$ and $H \cap Z(G) = G'$ by virtue of [9, Lemma 2.1]. Now, for the sake of clarity, we consider two cases.

Case1. First assume that G' lies in a central subgroup K of exponent p^2 . Therefore, one can check that there exists a central subgroup T such that $G = H \cdot K \times T \cong$

$E(2) \times T$. Thus, when T is an elementary abelian p -group by using [8, Theorem 2.2.10] and Theorem 4, we have

$$\begin{aligned} |\mathcal{M}(G)| &= |\mathcal{M}(E(2))||\mathcal{M}(T)||E(2)^{ab} \otimes T| \\ &= 2m^2 + m - 1 + \frac{1}{2}(n - 2m - 2)(n - 2m - 3) + 2m + 1(n - 2m - 2) \\ &= \frac{1}{2}(n - 1)(n - 2) - 1. \end{aligned}$$

In the case T is not elementary abelian, a similar method and [9, Lemma 2.2] asserts that

$$|\mathcal{M}(G)| \leq p^{\frac{1}{2}(n^2 - 5n + 4)} \leq p^{\frac{1}{2}(n-1)(n-2)-2}.$$

Case2. G' has a complement T in $Z(G)$, and hence $G = H \times T$ where T is not elementary abelian, and so by invoking [9, Lemma 2.2] and [8, Theorems 2.2.10 and 3.3.6], $|\mathcal{M}(G)| \leq p^{\frac{1}{2}(n-1)(n-2)-2}$. $\square$

Theorem 6. *Let G be a p -group of order p^n , G^{ab} be elementary abelian of order p^{n-1} and $Z(G)$ be of exponent p . Then G has the property $s(G) = 2$ if and only if it is isomorphic to one of the following groups.*

$$Q_8 \times \mathbb{Z}_2^{(n-3)}, E_2 \times \mathbb{Z}_p^{(n-3)} \text{ or } H \times \mathbb{Z}_p^{(n-2m-1)},$$

where H is extra special of order p^{2m+1} $m \geq 2$.

Proof. It is obtained via Theorem 4, [8, Theorems 2.2.10 and 3.3.6] and assumption. $\square$

Lemma 7. *Let G be a p -groups of order p^4 and $|G'| = p$. Then G has the property $s(G) = 2$ if and only if G is isomorphic to the one of the following groups.*

- (1) $Q_8 \times \mathbb{Z}_2$,
- (2) $\langle a, b \mid a^4 = 1, b^4 = 1, [a, b, a] = [a, b, b] = 1, [a, b] = a^2b^2 \rangle$
- (2) $\langle a, b, c \mid a^2 = b^2 = c^2 = 1, abc = bca = cab \rangle$.
- (4) $E_4 \cong E_1(2)$,
- (5) $E_2 \times \mathbb{Z}_p$,
- (6) $\langle a, b \mid a^{p^2} = 1, b^p = 1, [a, b, a] = [a, b, b] = 1 \rangle$,

Proof. It is obtained by using Theorems 4, 6 and a result of [10, Lemma 3.5]. $\square$

The structure of all p -group of order p^n is characterized with the property $s(G) = 2$ and $|G'| = p$. Now, we may suppose that $|G'| = p^2$.

Lemma 8. *There is no p -group of order p^n ($n \geq 5$) with $s(G) = 2$, where $|G'| = p^2$ and $G' \not\subseteq Z(G)$.*

Proof. First assume that $|Z(G)| = p^2$, since $Z(G)$ is elementary by Lemma 4. Let K be a central subgroup of order p , such that $|(G/K)'| = p^2$. It is seen that

$$|\mathcal{M}(G)| \leq |\mathcal{M}(G/K)||K \otimes G/(K \times G')| \leq |\mathcal{M}(G/K)| p^{n-3}$$

by [7, Theorem 4.1]. On the other hand, [9, 12, Main Thoerems] imply that $|\mathcal{M}(G/K)| \leq p^{\frac{1}{2}(n-2)(n-3)-1}$, and hence $|\mathcal{M}(G)| \leq p^{\frac{1}{2}(n-1)(n-2)-2}$.

In case $|Z(G)| = p^3$, there exists a central subgroup K of order p^2 such that $G' \cap K = 1$. The rest of proof is obtained similar to the pervious case.

When $|Z(G)| = p$, since G is nilpotent of class 3, the result is deduced by [8, Proposition 3.1.11]. $\square$

Theorem 9. *Let G be a p -group of order p^n ($n \geq 5$) and $|G'| = p^2$ with $s(G) = 2$. Then*

$$G \cong \mathbb{Z}_p \times (\mathbb{Z}_p^{(4)} \rtimes_{\theta} \mathbb{Z}_p) \quad (p \neq 2).$$

By the results of Lemmas 4 and 8, we may assume that $G' \subseteq Z(G)$ and $Z(G)$ is of exponent p . We consider three cases relative to $|Z(G)|$.

Case1. Assume that $|Z(G)| = p^4$, there exists a central subgroup K of order p^2 such that $K \cap G' = 1$. [9, Main Theorem] implies that $|\mathcal{M}(G/K)| \leq p^{\frac{1}{2}(n-3)(n-4)}$, and so $|\mathcal{M}(G)| \leq p^{\frac{1}{2}(n-1)(n-2)-1}$ due to [7, Theorem 4.1].

Case2. In the case $|Z(G)| = p^2$, we have $G' = Z(G)$. Moreover [12, Main Theorem] deduces that $n \geq 6$ and so there exists a central subgroup K such that $G/K \cong H \times Z(G/K)$ where H is a extra special p -groups of order p^{2m+1} $m \geq 2$, thus

$$|\mathcal{M}(G)| \leq p^{n-3} |\mathcal{M}(G/K)| \leq p^{n-3} p^{\frac{1}{2}(n-1)(n-4)} \leq p^{\frac{1}{2}(n-1)(n-2)-2}.$$

Case3. Now, we may assume that $|Z(G)| = p^3$. Let K be a complement of G' in $Z(G)$, so [11, Main Theorem] asserts that $|\mathcal{M}(G/K)| \leq p^{\frac{1}{2}(n-2)(n-3)}$. On the other hand, [7, Theorem 4.1] and assumption imply that

$$\begin{aligned} p^{\frac{1}{2}(n-1)(n-2)-1} = |\mathcal{M}(G)| &\leq |\mathcal{M}(G/K)| |K \otimes G/Z(G)| \\ &\leq |\mathcal{M}(G/K)| p^{n-3}, \end{aligned}$$

so we should have $|\mathcal{M}(G/K)| = p^{\frac{1}{2}(n-2)(n-3)}$ and $G/Z(G)$ is elementary abelian. Now, since $|\mathcal{M}(G/K)| = p^{\frac{1}{2}(n-2)(n-3)}$ and $|(G/K)'| = p^2$, by using [12, Main Theorem], $G/K \cong \mathbb{Z}_p^{(4)} \rtimes_{\theta} \mathbb{Z}_p$ ($p \neq 2$). Moreover, [3, Proposition 1] and assumption show that G^{ab} is elementary abelian. Hence, it is readily shown that

$$G \cong \mathbb{Z}_p \times (\mathbb{Z}_p^{(4)} \rtimes_{\theta} \mathbb{Z}_p) \quad (p \neq 2).$$

Theorem 10. *Let G be a group of order p^4 with $s(G) = 2$ and $|G'| = p^2$. Then G is isomorphic to the one of the following groups.*

- (1) $\langle a, b \mid a^9 = b^3 = 1, [a, b, a] = 1, [a, b, b] = a^6, [a, b, b, b] = 1 \rangle$,
- (2) $\langle a, b \mid a^p = 1, b^p = 1, [a, b, a] = [a, b, b, a] = [a, b, b, b] = 1 \rangle (p \neq 3)$.

Proof. The structure of these groups has been characterized in [10, Lemma 3.6]. $\square$

We summarize all results as follows

Theorem 11. *Let G be a group of order p^n . Then G has a property $s(G) = 2$ if and only if isomorphic to the one of the following groups.*

- (1) $E(2) \times \mathbb{Z}_p^{(n-2m-2)}$,
- (2) $E_2 \times \mathbb{Z}_p^{(n-3)}$,
- (3) $Q_8 \times \mathbb{Z}_2^{(n-3)}$,
- (4) $H \times \mathbb{Z}_p^{(n-2m-1)}$, where H is an extra special p -group of order p^{2m+1} ($m \geq 2$),
- (5) $\langle a, b \mid a^4 = 1, b^4 = 1, [a, b, a] = [a, b, b] = 1, [a, b] = a^2 b^2 \rangle$
- (6) $\langle a, b, c \mid a^2 = b^2 = c^2 = 1, abc = bca = cab \rangle$.
- (7) $\langle a, b \mid a^{p^2} = 1, b^p = 1, [a, b, a] = [a, b, b] = 1 \rangle$,
- (8) $\mathbb{Z}_p \times (\mathbb{Z}_p^{(4)} \rtimes_{\theta} \mathbb{Z}_p)$ ($p \neq 2$),
- (9) $\langle a, b \mid a^9 = b^3 = 1, [a, b, a] = 1, [a, b, b] = a^6, [a, b, b, b] = 1 \rangle$,
- (10) $\langle a, b \mid a^p = 1, b^p = 1, [a, b, a] = [a, b, b, a] = [a, b, b, b] = 1 \rangle (p \neq 3)$.

REFERENCES

- [1] Berkovich, Ya. G. (1991). On the order of the commutator subgroups and the Schur multiplier of a finite p -group. *J. Algebra* 144: 269-272.
- [2] Ellis, G. (1999). On the Schur multiplier of p -groups. *Comm. Algebra* 27(9): 4173-4177.
- [3] Ellis, G., Wiegold, J. (1999). A bound on the Schur multiplier of a prime-power group. *Bull. Austral. Math. Soc.* 60: 191-196.
- [4] Gaschütz, W., Neubüser, J., Yen, T. (1967). Über den Multiplikator von p -Gruppen, *Math. Z.* 100: 93-96.
- [5] Green, J.A., (1956). On the number of automorphisms of a finite group, *Proc. Roy. Soc. A* 237: 574-581.
- [6] Jones, M. R. (1972). Multiplicators of p -groups. *Math. Z.* 127: 165-166.
- [7] Jones, M.R. (1973). Some inequalities for the multiplier of a finite group. *Proc. Amer. Math. Soc.* 39 450-456.
- [8] G. Karpilovsky, The Schur multiplier, London Math. Soc. Monogr. (N.S.) 2 (1987).
- [9] P. Niroomand.(2009). On the order of Schur multiplier of non-abelian p -groups. *J. Algebra* 322: 4479-4482.
- [10] P. Niroomand. Characterizing finite p -groups by their Schur multipliers. Preprint.(2010) <http://arxiv.org/abs/1001.4256v1>.
- [11] P. Niroomand. Characterizing finite p -groups by their Schur multipliers, $t(G) = 5$. Preprint. (2010) <http://arxiv.org/abs/1001.4257v1>.
- [12] P. Niroomand. A note on the Schur multiplier of prime power groups. submitted.
- [13] Zhou, X. (1994). On the order of the Schur multiplier of finite p -groups. *Comm. Algebra* 22: 1-8.

E-mail address: `niroomand@dubs.ac.ir`

SCHOOL OF MATHEMATICS AND COMPUTER SCIENCE, DAMGHAN UNIVERSITY OF BASIC SCIENCES, DAMGHAN, IRAN